Noname manuscript No.
(will be inserted by the editor)

On a multiplicative version of Bloch's conjecture

Robert Laterveer

arXiv:1609.08798v1 [math.AG] 28 Sep 2016

Received: date / Accepted: date

Abstract A theorem of Esnault, Srinivas and Viehweg asserts that if the Chow group of 0–cycles of a smooth complete complex variety decomposes, then the top–degree coherent cohomology group decomposes similarly. In this note, we prove that (a weak version of) the converse holds for varieties of dimension at most 5 that have finite–dimensional motive and satisfy the Lefschetz standard conjecture. The proof is based on Vial's construction of a refined Chow–Künneth decomposition for these varieties.

Keywords Algebraic cycles · Chow groups · Intersection product · Finite–dimensional motives · Bloch–Beilinson conjectures

Mathematics Subject Classification (2010) 14C15 · 14C25 · 14C30

1 Introduction

Let X be a smooth complete variety of dimension n defined over $\mathbb{C}$. In the 1992 paper [9], Esnault, Srinivas and Viehweg study the multiplicative behaviour of the Chow ring A^*X versus the multiplicative behaviour of the cohomology of X. We now state the part of their result that is relevant to us. For a given partition $n = n_1 + \cdots + n_r$ (with $n_i \in \mathbb{N}_{>0}$), let us consider the following properties:

(P1) There exists a Zariski open $V \subset X$, such that intersection product induces a surjection

$$A^{n_1}V_{\mathbb{Q}} \otimes A^{n_2}V_{\mathbb{Q}} \otimes \cdots \otimes A^{n_r}V_{\mathbb{Q}} \ \to \ A^nV_{\mathbb{Q}} \ ;$$

(P2) There exists a Zariski open $V \subset X$, such that cup product induces a surjection

$$H^{n_1}(V,\mathbb{Q}) \otimes H^{n_2}(V,\mathbb{Q}) \otimes \cdots \otimes H^{n_r}(V,\mathbb{Q}) \ \to \ H^n(V,\mathbb{Q})/N^1$$

(here N^* denotes the coniveau filtration);
(P3) Cup product induces a surjection

$$H^{n_1}(X,\mathcal{O}_X) \otimes H^{n_2}(X,\mathcal{O}_X) \otimes \cdots \otimes H^{n_r}(X,\mathcal{O}_X) \ \to \ H^n(X,\mathcal{O}_X) \ .$$

In these terms, what Esnault, Srinivas and Viehweg prove is the following:

Theorem (Esnault–Srinivas–Viehweg [9]) *Let X be a smooth complete variety of dimension n over $\mathbb{C}$. Then (P1) implies (P3), and (P2) implies (P3).*

CNRS - IRMA, Université de Strasbourg
7 rue René Descartes
67084 Strasbourg cedex
France
E-mail: laterv@math.unistra.fr

The implication from (P1) to (P3) is a kind of Mumford theorem [25], and the proof in [9] is motivated by Bloch's proof [6], [5] of Mumford's theorem using a "decomposition of the diagonal" argument. As noted in [9, remark 2], the generalized Hodge conjecture would imply that (P2) and (P3) are equivalent.[1]

It seems natural to conjecture the converse implication (this is discussed in [9, remark 3]):

Conjecture *Let X be a smooth complete variety. Then (P2) implies (P1).*

This can be considered a multiplicative version of Bloch's conjecture; indeed for surfaces of geometric genus 0 this conjecture is equivalent to Bloch's conjecture [9, remark 3].

The object of this note is to show this conjecture can be proven in some special cases:

Theorem *Let X be a smooth projective variety of dimension $n \leq 3$, rationally dominated by curves. Then (P2) implies (P1).*

This follows from a more general statement (theorem 3). Actually, our argument works for varieties X of dimension up to 5 that satisfy the Lefschetz standard conjecture and have finite–dimensional motive in the sense of Kimura [21], provided we replace (P2) by a variant involving Vial's niveau filtration $\widetilde{N}^*$ [34] instead of N^*.

This result is hardly surprising: since the appearance of Kimura's landmark paper [21], where it is shown that finite–dimensionality implies the Bloch conjecture for surfaces, there have been a great many results attesting to the usefulness of finite–dimensionality [17], [1], [12], [18], [19], [10], [11], [34], [39], [24]. The present note is but one more instance of this general principle, illustrating how nicely finite–dimensionality allows to bridge the abyss separating homological equivalence from rational equivalence.

Conventions *In this note, the word variety will refer to an irreducible reduced scheme of finite type over $\mathbb{C}$, endowed with the Zariski topology. A subvariety is a (possibly reducible) reduced subscheme which is equidimensional. The Chow group of j–dimensional algebraic cycles on X with $\mathbb{Q}$–coefficients modulo rational equivalence is denoted A_jX; for X smooth of dimension n the notations A_jX and $A^{n-j}X$ will be used interchangeably. Caveat: note that what we denote A^jX is elsewhere often denoted $CH^j(X)_{\mathbb{Q}}$. The Griffiths group $Griff^j$ is the group of codimension j cycles that are homologically trivial modulo algebraic equivalence, again with $\mathbb{Q}$–coefficients. In an effort to lighten notation, we will often write H^jX or H_jX to indicate singular cohomology $H^j(X,\mathbb{Q})$ resp. Borel–Moore homology $H_j(X,\mathbb{Q})$.*

2 Preliminaries

Let X be a smooth projective variety of dimension n, and $h \in H^2(X,\mathbb{Q})$ the class of an ample line bundle. The hard Lefschetz theorem asserts that the map

$$L^{n-i}\colon H^i(X,\mathbb{Q}) \to H^{2n-i}(X,\mathbb{Q})$$

obtained by cupping with h^{n-i} is an isomorphism, for any $i < n$. One of the standard conjectures asserts that the inverse isomorphism is algebraic.

Definition 1 (Lefschetz standard conjecture) Given a variety X, we say that $B(X)$ holds if for all ample h, and all $i < n$ the isomorphism

$$(L^{n-i})^{-1}\colon H^{2n-i}(X,\mathbb{Q}) \stackrel{\cong}{\to} H^i(X,\mathbb{Q})$$

is induced by a correspondence.

Remark 1 It is known that $B(X)$ holds for the following varieties: curves, surfaces, abelian varieties [22], [23], 3folds not of general type [30], n–dimensional varieties X which have $A_i(X)$ supported on a subvariety of dimension $i+2$ for all $i \leq \frac{n-3}{2}$ [31, Theorem 7.1], n–dimensional varieties X which have $H_i(X) = N^{\llcorner \frac{i}{2} \lrcorner}H_i(X)$ for all $i > n$ [32, Theorem 4.2], products and hyperplane sections of any of these.

[1] It is somewhat frustrating that it is not known unconditionally whether (P1) implies (P2), i.e. without assuming the generalized Hodge conjecture. Apparently Esnault, Srinivas and Viehweg had claimed to prove this in an earlier version of their paper, but the argument was found to be incomplete [9, remark 2].

Definition 2 (Coniveau filtration [7]) Let X be a quasi–projective variety. The *coniveau filtration* on cohomology and on homology is defined as

$$N^c H^i(X,\mathbb{Q}) = \sum \mathrm{Im}\big(H^i_Y(X,\mathbb{Q}) \to H^i(X,\mathbb{Q})\big) \ ;$$
$$N^c H_i(X,\mathbb{Q}) = \sum \mathrm{Im}\big(H_i(Z,\mathbb{Q}) \to H_i(X,\mathbb{Q})\big) \ ,$$

where Y runs over codimension $\geq c$ subvarieties of X, and Z over dimension $\leq i-c$ subvarieties.

Vial introduced the following variant of the coniveau filtration:

Definition 3 (Niveau filtration [34]) Let X be a smooth projective variety. The *niveau filtration* on homology is defined as

$$\widetilde{N}^j H_i(X) = \sum_{\Gamma \in A_{i-j}(Z\times X)} \mathrm{Im}\big(H_{i-2j}(Z) \to H_i(X)\big) \ ,$$

where the union runs over all smooth projective varieties Z of dimension $i-2j$, and all correspondences $\Gamma \in A_{i-j}(Z\times X)$. The niveau filtration on cohomology is defined as

$$\widetilde{N}^c H^i X := \widetilde{N}^{c-i+n} H_{2n-i} X \ .$$

Remark 2 The niveau filtration is included in the coniveau filtration:

$$\widetilde{N}^j H^i(X) \subset N^j H^i(X) \ .$$

These two filtrations are expected to coincide; indeed, Vial shows this is true if and only if the Lefschetz standard conjecture is true for all varieties [34, Proposition 1.1].

The main ingredient we will use in this note is Kimura's nilpotence theorem:

Theorem 1 (Kimura [21]) *Let X be a smooth projective variety of dimension n with finite–dimensional motive. Let $\Gamma \in A^n(X\times X)$ be a correspondence which is numerically trivial. Then there is $N\in\mathbb{N}$ such that*

$$\Gamma^{\circ N} = 0 \quad \in A^n(X\times X) \ .$$

We refer to [21], [1], [12], [27] for the definition of finite–dimensional motive. Conjecturally, any variety has finite–dimensional motive [21]. What mainly concerns us in the scope of this note, is that there are quite a few non–trivial examples, giving rise to interesting applications:

Remark 3 The following varieties have finite–dimensional motive: abelian varieties, varieties dominated by products of curves [21], $K3$ surfaces with Picard number 19 or 20 [28], surfaces not of general type with vanishing geometric genus [10, Theorem 2.11], Godeaux surfaces [10], 3folds and 4folds with nef tangent bundle [13], [33, Example 3.16], [14], certain 3folds of general type [35, Section 8], varieties of dimension ≤ 3 rationally dominated by products of curves [33, Example 3.15], varieties X with $A^i_{AJ}X_{\mathbb{Q}}=0$ for all i [32, Theorem 4], products of varieties with finite–dimensional motive [21].

Remark 4 It is worth pointing out that up till now, all examples of finite-dimensional motives happen to be in the tensor subcategory generated by Chow motives of curves. On the other hand, "many" motives are known to lie outside this subcategory, e.g. the motive of a general hypersurface in $\mathbb{P}^3$ [3, Remark 2.34] (here "general" means "outside a countable union of Zariski–closed proper subsets").

There exists another nilpotence result, which predates and prefigures Kimura's theorem:

Theorem 2 (Voisin [37], Voevodsky [36]) *Let X be a smooth projective algebraic variety of dimension n, and $\Gamma \in A^n(X\times X)$ a correspondence which is algebraically trivial. Then there is $N\in\mathbb{N}$ such that*

$$\Gamma^{\circ N} = 0 \quad \in A^n(X\times X) \ .$$

3 Main result

In this section, we prove the main result of this note:

Theorem 3 *Let X be a smooth projective variety over $\mathbb{C}$ of dimension n. Suppose*
(i) $n \leq 5$;
(ii) $B(X)$ is true;
(iii) X has finite–dimensional motive, or Griff$^n(X \times X) = 0$.
Given $n_i \in \mathbb{N}$ with $n = n_1 + \cdots n_r$, suppose that
(P2) Cup product induces a surjection

$$H^{n_1}(X) \otimes H^{n_2}(X) \otimes \cdots \otimes H^{n_r}(X) \ \to \ H^n(X)/\widetilde{N}^1 \ .$$

Then
(P1) There exists an open $V \subset X$, such that intersection product induces a surjection

$$A^{n_1}V \otimes A^{n_2}V \otimes \cdots \otimes A^{n_r}V \ \to \ A^nV \ .$$

In dimension ≤ 3, the niveau filtration $\widetilde{N}^1$ can be replaced by the coniveau filtration N^1, and we obtain the result announced in the introduction:

Corollary 1 *Let X be as in theorem 3, and of dimension $n \leq 3$. Then condition (P2) can be replaced by*
(P2′) There is an open $V \subset X$ such that cup product induces a surjection

$$H^{n_1}(V) \otimes \cdots \otimes H^{n_r}(V) \ \to \ H^n(V)/N^1 \ .$$

If $n = 2$, condition (P2) can be replaced by
(P2″) There is an open $V \subset X$ such that cup product induces a surjection

$$H^1(V) \otimes H^1(V) \ \to \ H^2(V)/F^1 \ ,$$

where F^* denotes the Hodge filtration.

Proof (of corollary 1) It is easily seen that (P2′) implies surjectivity of

$$H^{n_1}(X) \otimes \cdots \otimes H^{n_r}(X) \ \to \ H^n(X)/N^1 \ .$$

Suppose now $n = 3$. Since

$$N^1H^3X = \widetilde{N}^1H^3X$$

[34, page 415 "Properties"], the above is equivalent to condition (P2) of theorem 3. The case $n = 2$ is similar, using that

$$\widetilde{N}^1H^2X = N^1H^2X = F^1H^2X \ .$$

Proof (of theorem 3)

Since X satisfies $B(X)$, the Künneth components π_i are algebraic [22], [23]. Because the dimension is at most 5, the variety X satisfies conditions (*) and (**) of Vial's [34]. This implies the existence of a refined Chow–Künneth decomposition of the diagonal

$$\Delta = \sum_{i,j} \Pi_{i,j} \quad \in A^n(X \times X)$$

(loc. cit., Theorems 1 and 2). Here the $\Pi_{i,j}$ are mutually orthogonal idempotents, which act on homology as projectors

$$(\Pi_{i,j})_* : \ \ H_*(X) \ \to \ \mathrm{Gr}^j_{\widetilde{N}} H_i(X) \to H_*(X) \ .$$

(Note that $\mathrm{Gr}^j_{\widetilde{N}} H_i(X)$ is a priori not a subspace of $H_i(X)$; however, over $\mathbb{C}$ the existence of a polarization gives a canonical identification with a subspace of $H_i(X)$ [34].) Assumption (P2) translates into the fact that the morphism of homological motives

$$f = \Pi_{n,0} \circ \Delta^r \circ (\pi_{n_1} \times \cdots \times \pi_{n_r}) : \ \ (X, \pi_{n_1}) \otimes \cdots \otimes (X, \pi_{n_r}) \ \to \ (X, \Pi_{n,0}) \ \ \in \mathcal{M}_{hom}$$

is surjective, where Δ^r is the class in $A^{nr}(X^{r+1})$ of the "small diagonal"

$$\Delta^r := \{(x,x,\ldots,x), x\in X\} \ \subset X^{r+1}\ .$$

To see this, note that since $B(X)$ holds and we are in characteristic 0, homological and numerical equivalence coincide on X and its powers [22], [23]. Thus, using Jannsen's semisimplicity result [15], the motives (X,π_i) and $(X,\Pi_{n,0})$ are contained in a full semisimple abelian subcategory $\mathcal{M}_0$ of the category $\mathcal{M}_{hom}$ of motives with respect to homological equivalence (for $\mathcal{M}_0\subset\mathcal{M}_{hom}$ one can take the subcategory generated by varieties that are known to satisfy the Lefschetz standard conjecture). Hence, we get a decomposition

$$(X,\Pi_{n,0}) = \mathrm{Im} f\oplus M' \ \in \mathcal{M}_0\subset\mathcal{M}_{hom}\ .$$

But assumption (P2) gives that

$$H^*(M')=0\ ,$$

so that $M'=0\in\mathcal{M}_{hom}$.

By semisimplicity of $\mathcal{M}_0$, the surjection

$$f=\Pi_{n,0}\circ\Delta^r\circ(\pi_{n_1}\times\cdots\times\pi_{n_r})\colon\ \ (X,\pi_{n_1})\otimes\cdots\otimes(X,\pi_{n_r})\ \to\ (X,\Pi_{n,0})\ \in\mathcal{M}_{hom}$$

is a split surjection. That is, there exists a correspondence $C\in A^n(X\times X^r)$ such that

$$\Pi_{n,0}=\Pi_{n,0}\circ\Delta^r\circ(\pi_{n_1}\times\cdots\times\pi_{n_r})\circ C\ \in H^{2n}(X\times X)\ .$$

For brevity, we will henceforth write

$$\Delta_{(n_1,\ldots,n_r)}:=\Pi_{n,0}\circ\Delta^r\circ(\pi_{n_1}\times\cdots\times\pi_{n_r})\circ C\ \in A^n(X\times X)\ ,$$

where we suppose we have made the following choice for the Künneth components π_{n_i} modulo rational equivalence:

$$\pi_{n_i}=\sum_j \Pi_{n_i,j}\ \in A^n(X\times X)\ .$$

Since by construction [34, Theorems 1 and 2], all the $\Pi_{i,j}$ for $(i,j)\neq(n,0)$ are supported on $(D\times X)\cup(X\times D)$, for some divisor $D\subset X$, we get an equality modulo homological equivalence

$$\Delta=\Pi_{n,0}+\sum_{(i,j)\neq(n,0)}\Pi_{i,j}=\Delta_{(n_1,\ldots,n_r)}+\Gamma_1+\Gamma_2\ \in H^{2n}(X\times X)\ ,$$

with Γ_1,Γ_2 supported on $D\times X$ (resp. on $X\times D$).

Using one of the two nilpotence theorems (theorem 1 in case X has finite–dimensional motive, theorem 2 in case the Griffiths group vanishes), it follows there exists $N\in\mathbb{N}$ such that

$$\Big(\Delta-\Delta_{(n_1,\ldots,n_r)}-\Gamma_1-\Gamma_2\Big)^{\circ N}=0\ \in A^n(X\times X)\ .$$

Developing this expression gives

$$\Delta=\sum_k Q_k\ \in A^n(X\times X)\ ,$$

where each Q_k is a composition of $\Delta_{(n_1,\ldots,n_r)}$ and Γ_1 and Γ_2:

$$Q_k=Q_k^0\circ Q_k^1\circ\cdots\circ Q_k^{N'}\ \in A^n(X\times X)\ ,$$

for $Q_k^i\in\{\Delta_{(n_1,\ldots,n_r)},\Gamma_1,\Gamma_2\}$.

For reasons of dimension, Q_k does not act on A^nX as soon as Q_k contains at least one copy of Γ_1. It follows that

$$A^nX=\Delta_*A^nX=(\Delta_{(n_1,\ldots,n_r)}\circ(\text{something}))_*A^nX+(\Gamma_2\circ(\text{something}))_*A^nX\ . \qquad (1)$$

It is convenient to rewrite this as follows: define

$$\Delta'_{(n_1,\ldots,n_r)} := \Delta^r \circ (\pi_{n_1} \times \cdots \times \pi_{n_r}) \circ C \ \ \in A^n(X \times X) \ ,$$

so that

$$\Delta_{(n_1,\ldots,n_r)} = \Pi_{n,0} \circ \Delta'_{(n_1,\ldots,n_r)} \ \ \in A^n(X \times X) \ .$$

Since all the $\Pi_{i,j}$ with $(i,j) \neq (n,0)$ are supported on $(D \times X) \cup (X \times D)$ [34, Theorems 1 and 2], we have an equality

$$\Delta_{(n_1,\ldots,n_r)} = (\Delta - \sum_{(i,j)\neq(n,0)} \Pi_{i,j}) \circ \Delta'_{(n_1,\ldots,n_r)} = \Delta'_{(n_1,\ldots,n_r)} + \Gamma_1 + \Gamma_2 \ \ \in A^n(X \times X) \ ,$$

with Γ_1, Γ_2 supported on $D \times X$ resp. on $X \times D$. Now, equation (1) can be rewritten as

$$A^n X = (\Delta'_{(n_1,\ldots,n_r)} \circ (\text{something}))_* A^n X + (\Gamma_2 \circ (\text{something}))_* A^n X \ .$$

The first term decomposes (lemma 1 below), and the second term is supported on D; this proves the theorem with $V = X \setminus D$.

Lemma 1 *Set–up as above. Then*

$$(\Delta'_{(n_1,\ldots,n_r)})_* A^n X \subset \mathit{Im}\big(A^{n_1}X \otimes \cdots \otimes A^{n_r}X \ \stackrel{\iota}{\to} \ A^n X\big) \ ,$$

where ι denotes intersection product.

Proof Recall that by definition,

$$\Delta'_{(n_1,\ldots,n_r)} = \Delta^r \circ (\pi_{n_1} \times \cdots \times \pi_{n_r}) \circ C \ \ \in A^n(X \times X) \ .$$

The point is that the π_{n_i} are supported on $Y_{n_i} \times X$, for some $Y_{n_i} \subset X$ of dimension n_i, i.e. there exist correspondences $\pi'_{n_i} \in A_n(Y_{n_i} \times X)$ pushing forward to π_{n_i} [34, Theorems 1 and 2] (actually, this can even be achieved with Y_{n_i} a smooth hyperplane section of dimension n_i [19, Theorem 7.7.4]). The action of the correspondence $\Delta'_{(n_1,\ldots,n_r)}$ on 0–cycles thus factors as

$$\begin{array}{ccccccc}
A^n X & & \xrightarrow{(\Delta'_{(n_1,\ldots,n_r)})_*} & & & & A^n X \\
\downarrow = & & & & & & \uparrow = \\
A^n X & \xrightarrow{C_*} & A^n(X^r) & \xrightarrow{(\pi_{n_1}\times\cdots\times\pi_{n_r})_*} & A^n(X^r) & \xrightarrow{(\Delta^r)_*} & A^n X \\
& & \downarrow & & \uparrow = & & \uparrow = \\
& & A^n(Y_{n_1}\times\cdots\times Y_{n_r}) & \xrightarrow{(\pi'_{n_1}\times\cdots\times\pi'_{n_r})_*} & A^n(X^r) & \xrightarrow{(\Delta^r)_*} & A^n X \\
& & \uparrow \times_r & & \uparrow & & \uparrow = \\
& & A^{n_1}(Y_{n_1})\otimes\cdots\otimes A^{n_r}(Y_{n_r}) & \xrightarrow{(\pi'_{n_1})_*\otimes\cdots\otimes(\pi'_{n_r})_*} & A^{n_1}(X)\otimes\cdots\otimes A^{n_r}(X) & \stackrel{\iota}{\to} & A^n X
\end{array}$$

The arrow labelled $\times_r$ is surjective (because it is a map on 0–cycles); since the diagram commutes this implies the lemma.

Remark 5 Here are some non–trivial cases where property (P1) is known to hold: abelian varieties (with $V = X$ and all the $n_j = 1$, cf. [4]); the variety of lines of a cubic threefold (with $V = X$ and $n_1 = n_2 = 1$, cf. [6, Example 1.7]); the variety of lines of a cubic fourfold (with $V = X$ and $n_1 = n_2 = 2$, cf. [29, Theorem 20.2]). Note that in the last two cases, finite–dimensionality of the motive is not known, so (P1) can *not* be deduced from theorem 3; one needs some geometric arguments to establish (P1).

Remark 6 The assumption "Griff$^n(X\times X)=0$" in theorem 3 is mainly of theoretical interest, and not practically useful. Indeed, there are precise conjectures (based on the Bloch–Beilinson conjectures) saying how the coniveau filtration on cohomology should influence Griffiths groups [16]. For $n=2$, it is conjectured that if $H^1(X)=0$ then Griff$^2(X\times X)=0$. For $n=3$, it is conjectured that if $h^{0,2}(X)=h^{0,3}(X)=0$ then Griff$^3(X\times X)=0$. For $n=4$, if $h^{2,0}(X)=h^{3,0}(X)=h^{4,0}(X)=h^{2,1}(X)=0$ then Griff$^4(X\times X)$ should vanish. These predictions are particular instances of [16, Corollary 6.8].

Unfortunately, it seems these conjectures are not known in any non–trivial cases (i.e., outside of the range of varieties with Abel–Jacobi trivial Chow groups); it would be very interesting to find (non–trivial) examples where they can be proven !

Remark 7 The Chow motive $(X,\Pi_{n,0})$ (which for varieties verifying (*) and (**) of [34] is unique up to isomorphism) can be considered the "most transcendental part" of the motive (X,Δ). When X is a surface, $(X,\Pi_{2,0})$ is the transcendental part denoted $t_2(X)$ (and studied in detail) in [19].

Actually, following [19] one might hope that $\Pi_{n,0}$ can be linked with the theory of birational motives of Kahn–Sujatha [20]; this would perhaps give a more conceptual proof (or an extension ?) of theorem 3. I have not looked into this yet.

Remark 8 The argument of theorem 3 also shows the following: suppose the standard conjecture of Lefschetz type holds universally. Then for any variety with finite–dimensional motive, (P2) implies (P1).

Remark 9 Suppose X satisfies the hypotheses of theorem 3, so that

$$A^{n_1}V\otimes A^{n_2}V\otimes\cdots\otimes A^{n_r}V\ \to\ A^nV$$

is surjective, for some open $V\subset X$. It seems interesting to ask how many simple tensors are needed to generate A^nV. For a given $n_1,\ldots,n_r$, let's say $a\in A^nV$ is *k–decomposable* if there is an expression

$$a=\sum_{j=1}^k a^1_j\cdot a^2_j\cdots a^r_j\ \ \in A^nV\ ,$$

with $a^i_j\in A^{n_i}V$. This is related to unpublished work of Nori, discussed in [9, remark 5]. According to loc. cit., Nori proves that for any X with $H^n(X,\mathcal{O}_X)\neq 0$ and any k, there exist elements in A^nX that are not k–decomposable (with respect to any $(n_1,\ldots,n_r)$ with $r>1$).[2] It seems likely the same is true for A^nV.

The only thing I am able to prove is the following: for X satisfying the hypotheses of theorem 3, there is an open $V\subset X$ such that each point $v\in V$ is 1–decomposable in A^nV. (To see this, one uses (P1) to obtain a Bloch–Srinivas style decomposition of the diagonal

$$\Delta=C_1\cdot C_2\cdots C_r+\Gamma_1+\Gamma_2\ \ \in A^n(X\times X)\ ,$$

with $C_i\in A^{n_i}(X\times X)$ and Γ_j as before; this is done in [9] using the method of [8]. Given a point $v\in V$, let τ_v denote the inclusion $v\times X\hookrightarrow X\times X$, and let C_i^v denote the restriction $C_i^v=(\tau_v)^*(C_i)\in A^{n_i}(v\times X)$. Now, note that

$$\begin{aligned} v=\Delta_*v=(C_1\cdots C_r)_*v&=(p_2)_*\big((v\times X)\cdot C_1\cdots C_r\big)\\ &=(p_2)_*\big((\tau_v)_*(C_1^v\cdots C_r^v)\big)\\ &=C_1^v\cdots C_r^v\ \ \in A^nV\ .)\end{aligned}$$

For general 0–cycles on the other hand, even when (as in the case of theorem 3) all 0–cycles are k–decomposable for some k, it seems unlikely k can be bounded.

Acknowledgements This note was stimulated by the Strasbourg 2014—2015 "groupe de travail" based on the monograph [38]. I want to thank all the participants of this groupe de travail for the very pleasant and stimulating atmosphere, and their interesting lectures. Thanks to Charles Vial and the referee for helpful comments. Many thanks to Yasuyo, Kai and Len for providing excellent working conditions at home in Schiltigheim.

[2] To be precise, Nori's result is more general, as the notion of k–decomposability in [9] is broader than the notion discussed here: in loc. cit., an element $a\in A^nV$ is defined to be k–decomposable if it can be written as $a=\sum_{j=1}^k a_j\cdot b_j$ with $a_j\cdot b_j$ homogeneous of degree n.

References

1. Y. André, Motifs de dimension finie (d'après S.-I. Kimura, P. O'Sullivan...), Séminaire Bourbaki. Vol. 2003/2004. Astérisque 299 Exp. No. 929, viii, 115—145,
2. D. Arapura, Varieties with very little transcendental cohomology, in: Motives and algebraic cycles (R. de Jeu and J. Lewis, eds.), Fields Institute Communications, Amer. Math. Soc., Providence 2009,
3. J. Ayoub, Motives and algebraic cycles: a selection of conjectures and open questions, preprint,
4. S. Bloch, Some elementary theorems about algebraic cycles on abelian varieties, Invent. Math. 37 (1976), 215—228,
5. S. Bloch, On an argument of Mumford in the theory of algebraic cycles, in: Géométrie algébrique (A. Beauville, ed.), Sijthoff and Noordhoff, Angers 1979,
6. S. Bloch, Lectures on algebraic cycles, Duke Univ. Press Durham 1980,
7. S. Bloch and A. Ogus, Gersten's conjecture and the homology of schemes, Ann. Sci. Ecole Norm. Sup. 4 (1974), 181—202,
8. S. Bloch and V. Srinivas, Remarks on correspondences and algebraic cycles, American Journal of Mathematics Vol. 105, No 5 (1983), 1235—1253,
9. H. Esnault, V. Srinivas and E. Viehweg, Decomposability of Chow groups implies decomposability of cohomology, in :"Journées de Géométrie Algébrique d'Orsay, Juillet 1992", Astérisque 218 (1993), 227—242,
10. V. Guletskiĭ and C. Pedrini, The Chow motive of the Godeaux surface, in: Algebraic Geometry, a volume in memory of Paolo Francia (M.C. Beltrametti, F. Catanese, C. Ciliberto, A. Lanteri and C. Pedrini, editors), Walter de Gruyter, Berlin New York, 2002,
11. V. Guletskiĭ and C. Pedrini, Finite–dimensional motives and the conjectures of Beilinson and Murre, K–Theory 30(3) (2003), 243—263,
12. F. Ivorra, Finite dimensional motives and applications (following S.-I. Kimura, P. O'Sullivan and others), in: Autour des motifs, Asian-French summer school on algebraic geometry and number theory, Volume III, Panoramas et synthèses, Société mathématique de France 2011,
13. J. Iyer, Murre's conjectures and explicit Chow–Künneth projectors for varieties with a nef tangent bundle, Transactions of the Amer. Math. Soc. 361 (2008), 1667—1681,
14. J. Iyer, Absolute Chow–Künneth decomposition for rational homogeneous bundles and for log homogeneous varieties, Michigan Math. Journal Vol.60, 1 (2011), 79—91,
15. U. Jannsen, Motives, numerical equivalence, and semi-simplicity, Invent. Math. 107(3) (1992), 447—452,
16. U. Jannsen, Equivalence relations on algebraic cycles, in: The arithmetic and geometry of algebraic cycles (B. Gordon et alii, eds.), Banff Conf. 1998, Kluwer,
17. U. Jannsen, On finite–dimensional motives and Murre's conjecture, in: Algebraic cycles and motives, volume 344 of London Math. Soc. Lecture Note Ser., Cambridge Univ. Press, Cambridge 2007,
18. B. Kahn, Equivalences rationnelle et numérique sur certaines variétés de type abélien sur un corps fini, Ann. Sci. Ec. Norm. Sup. 36 (2003), 977—1002,
19. B. Kahn, J. P. Murre and C. Pedrini, On the transcendental part of the motive of a surface, in: Algebraic cycles and motives. Vol. 2, volume 344 of London Math. Soc. Lecture Note Ser., 143—202, Cambridge Univ. Press, Cambridge 2007,
20. B. Kahn and R. Sujatha, Birational motives, I: pure birational motives, arXiv:0902.4902v4,
21. S. Kimura, Chow groups are finite dimensional, in some sense, Math. Ann. 331 (2005), 173—201,
22. S. Kleiman, Algebraic cycles and the Weil conjectures, in: Dix exposés sur la cohomologie des schémas, North–Holland Amsterdam, 1968, 359—386,
23. S. Kleiman, The standard conjectures, in: Motives (U. Jannsen et alii, eds.), Proceedings of Symposia in Pure Mathematics Vol. 55 (1994), Part 1,
24. R. Laterveer, A brief note concerning hard Lefschetz for Chow groups, to appear in Canadian Math. Bulletin, http://dx.doi.org/10.4153/CMB-2015-046-x,
25. D. Mumford, Rational equivalence of 0–cycles on surfaces, J. Math. Kyoto Univ. Vol. 9 No 2 (1969), 195—204,
26. J. Murre, On a conjectural filtration on the Chow groups of an algebraic variety, Indag. Math. (N.S.) 4 (1993), 177—201,
27. J. Murre, J. Nagel and C. Peters, Lectures on the theory of pure motives, Amer. Math. Soc. University Lecture Series 61, 2013,
28. C. Pedrini, On the finite dimensionality of a $K3$ surface, Manuscripta Mathematica 138 (2012), 59—72,
29. M. Shen and C. Vial, The Fourier transform for certain hyperKähler fourfolds, Memoirs of the AMS 240 (2016), no.1139,
30. S. Tankeev, On the standard conjecture of Lefschetz type for complex projective threefolds. II, Izvestiya Math. 75:5 (2011), 1047—1062,
31. C. Vial, Algebraic cycles and fibrations, Documenta Math. 18 (2013), 1521—1553,
32. C. Vial, Projectors on the intermediate algebraic Jacobians, New York J. Math. 19 (2013), 793—822,
33. C. Vial, Remarks on motives of abelian type, arXiv:1112.1080v2,
34. C. Vial, Niveau and coniveau filtrations on cohomology groups and Chow groups, Proceedings of the LMS 106(2) (2013), 410—444,
35. C. Vial, Chow-Künneth decomposition for 3- and 4-folds fibred by varieties with trivial Chow group of zero-cycles, J. Algebraic Geom. 24 (2015), 51—80,
36. V. Voevodsky, A nilpotence theorem for cycles algebraically equivalent to zero, Internat. Math. Research Notices 4 (1995), 187—198,
37. C. Voisin, Remarks on zero–cycles on self–products of varieties, in: Moduli of vector bundles (Proceedings of the Taniguchi Congress), Maruyama Ed., Decker 1994,
38. C. Voisin, Chow Rings, Decomposition of the Diagonal, and the Topology of Families, Princeton University Press, Princeton and Oxford, 2014,
39. Q. Yin, Finite–dimensionality and cycles on powers of K3 surfaces, Comment. Math. Helv. 90 (2015), 503—511.